

Substance	Grade
(3,3,3-Trifluoropropyl)chlorodimethylsilane	95%
(3,3,3-Trifluoropropyl)trichlorsilan	>97%
(3,3,3-Trifluoropropyl)trimethoxysilan	>97%
1,10-Phenanthroline	(anhydrous) for synthesis
1-Butanol	1-Butanol for analysis EMSURE®
1-Butanol	CPC fill bottle, working fluid
1H,1H,2H,2H-Perfluoro-1-octanol	97%
1H,1H,2H,2H-Perfluorooctyltriethoxysilan	> 97%
1-Propanol	ROTIPURAN® ≥99,5 %, p.a.
2,2-DMP; 2,2-dimethoxypropane	98%
2,6-Di-tert-butyl-p-cresol, BHT	for synthesis
2-Butanol	2-Butanol for analysis EMSURE®
2-Dodeceny succinic acid anhydride	pract.
2-Mercaptoethanol	≥99.0%
2-Methyl-2,4-pentandiol	≥99 %, for synthesis
2-propanol	99,50%
2-Propanol	99,50%
2-Propanol	99,50%
2-Propanol	99,50%
2-Propanol	99,50%
2-propanol	HiPerSolv, >99.9%
2-propanol	HiPerSolv, >99.9%
2-Propanol, isopropanol	ROTIPURAN® ≥99,8 %, p.a., ACS, ISO
2-Propanol, isopropanol	99,50%
2-Propanol, isopropanol	99,50%
2xYT Broth	for molecular biology
Acetate buffer (50mM Na+ / pH 3.5)	
Acetate buffer (50mM/pH 3.5)	
Acetic acid	ROTIPURAN® 100 %, p.a.
Acetic acid	ROTIPURAN® 100 %, p.a.
Acetic acid 100 %	ROTIPURAN® 100 %, p.a.
Aceton	für die Spektroskopie Uvasol®
Acetone	Emplura, 99,5%
Acetone	99,50%
Acetone	ROTIPURAN® ≥99,8 %, p.a., ACS, ISO
Acetone	Emplura, 99,5%
Acetone	EMPLURA®, Reagent Grade for synthesis, Supelco®
Acetone	EMPLURA®, Reagent Grade for synthesis, Supelco®
Acetonitrile	ROTISOLV® ≥99,8 %, for preparative HPLC
Acetonitrile	ACROS, Extra dry
Acetonitrile	ACROS, Extra dry
Acetonitrile	anhydrous (max. 0.003% H ₂ O) ≥99.95%, HiPerSolv CHROMANORM® Reag. Ph. Eur., USP, ACS, super gradient grade suitable for UPLC/UHPLC instruments
Acrylamide/Bis stock sol. 29:1 (40% w/v)	Rotiphorese® Gel 40 (29:1)
AEBSF hydrochloride	≥98 %, for biochemistry
Agar	for microbiology

Agar	Agar-Agar, bacteriological for bacteriology, extremely pure quality
Agar	Agar-Agar, Kobe I Kobe I, powdered, for microbiology
Agarose	Ultra-low Gelling Temperature, molecular biology grade
Agarose Standard	Roti®garose for DNA/RNA-electrophoresis
alpha Lactose Monohydrate	for microbiology
Ammonium acetate	BioXtra, ≥98%
Ammonium chloride	≥99,7 %, p.a., Ultra Quality
Ammonium dihydrogen phosphate	cryst. extra pure.
Ammonium formate	≥95 %, extra pure
Ammonium iron(II) sulphate hexahydrate	≥99 %, p.a., ISO
Ammonium nitrate	for analysis EMSURE® ACS.
Ammonium peroxodisulphate (APS)	≥98 %, p.a., ACS
Ammonium sulfite monohydrate	92%
Ammonium sulphate	Ammoniumsulfat min. 99,5 %
Ammonium sulphate	≥99%, GPR RECTAPUR®
Ammonium sulphate	≥99 %, p.a., ISO
Ammonium sulphate	≥99,5 %, p.a., ACS, ISO
Ammonium sulphate; not yet delivered	≥99.5%, Ultra Pure Grade
Ampicillin sodium Salt	≥97 %, BioScience-Grade, for molecular biology and biochemistry
ATP Dinatriumsalz	≥98 %, for biochemistry
Benzene	≥99% ACS, Riedel-de Haën™
BES	≥99.0% (titration)
BICINE	PUFFERAN® ≥99 %
BIS-TRIS	PUFFERAN® ≥99 %
Bluo-gal	
Boric Acid	BioXtra, ≥99.5%
Bovine Serum Albumin (BSA)	heat shock fraction, pH 7, ≥98%
Bromophenol blue indicator ACS	Bromophenol blue indicator ACS, Reag. Ph Eur.
Cacodylic acid sodium salt trihydrate; Sodium cacodylate	PUFFERAN® ≥98 %
Calcium acetate hydrate	BioXtra, ≥99% (titration)
Calcium carbonate	≥98,5 %, powdered
Calcium chloride dihydrate	BioUltra, for molecular biology, ≥99.5% (KT)
Calcium chloride dihydrate	
Calcium hydroxide	≥95 %, Ph.Eur., USP, BP
Calcium nitrate tetrahydrate	≥99 %, p.a., ACS
Calcium sulfate dihydrate	Calcium sulfate dihydrate precipitated for analysis EMSURE®.
CAPS	PUFFERAN® ≥99 %
CARBOXYMETHYLCELLULOSE SODIUM SALT	CARBOXYMETHYLCELLULOSE SODIUM SALT
Casamino acids	BD Bacto™ Casamino Acids
Cerium(IV) sulfate	Tetrahydrat, EMSURE® zur Analyse
Chloramphenicol	(R,R)-Chloramphenicol, 99+%
Chloroform	anhydrous, contains amylenes as stabilizer, ≥99%
Chlorotrimethylsilane	

Citric acid	monohydrate
Citric acid	monohydrate ≥98% for decalcifying
Citric acid	ACS reagent, ≥99.5%
Citric acid monohydrate	ACS reagent, ≥99.0%
Copper sulphate	
D(+) Sucrose / Saccharose	≥99,5 %, p.a. For microbiology and molecular biology.
D(+)-Galactose	≥98 %, for biochemistry
D(+)-Glucose	p.a., ACS, anhydrous
dAMP / 2'-Deoxyadenosine 5'-monophosphate	Sigma Grade, 98-100%
di-Ammonium Hydrogen Phosphate	≥98 %, p.a., ACS
dichloromethane	>99,8% (max.0,001% H2O, stab.: 2-Methyl-2-buten, 20 ppm)
Dichloromethane	for spectroscopy Uvasol®. CAS 75-09-2, chemical formula CH ₂ Cl ₂ ., for spectroscopy Uvasol®
dichloromethane	for spectroscopy Uvasol®. CAS 75-09-2, chemical formula CH ₂ Cl ₂ ., for spectroscopy Uvasol®
Diethyl ether	EMPLURA, stabilized with BHT/Ionol 15 ppm
Diethyl ether	EMPLURA, stabilized with BHT/Ionol 15 ppm
Diethylene glycol monobutyl ether	Sigma-Aldrich®
Difco™ Nutrient Broth	BD Difco™ Nutrient Broth
Dimethyl sulfoxide / DMSO	for molecular biology
Dimethylsulfoxid	
di-Potassium hydrogen phosphate	≥99 %, p.a., anhydrous
di-Potassium hydrogen phosphate	anhydrous ≥99%, puriss. p.a. ACS, Fluka™
Disodium hydrogen phosphate	≥98 %, Ph.Eur., USP
di-Sodium hydrogen phosphate heptahydrate / Sodium phosphate dibasic	≥98 %, p.a., ACS
di-Sodium tartrate dihydrate	≥99 %, p.a.
D-Mannitol	≥98%
DMSO	≥99,5 %, BioScience-Grade, for molecular biology
D-Sorbitol / D-Glucitol	≥98%
DTT (1,4-Dithiothreit)	≥99 %, p.a.
EDTA (Ethylenediamine tetraacetic acid tetrasodium salt hydrate)	86-88%
EDTA, Ethylenediamine tetraacetic acid	≥99 %, p.a., ACS. For molecular biology and biochemistry.
EGTA	≥99 %, p.a. For molecular biology and biochemistry.
Erythromycin	Cell Culture Grade
ESF 921 Media	
Ethanol	99,80%
Ethanol	Absolut
Ethanol	ROTIPURAN® ≥99,8 %, p.a.
Ethanol	Absolut, Emplura
Ethanol	Absolut, Emplura
Ethanol	Absolut, Emplura, "degased"
Ethanol	70%
Ethanol denatured	≥99,8 %, with ca. 1 % MEK

Ethanol denatured	≥99,8 %, with ca. 1 % MEK
Ethanolamine	for synthesis, Sigma-Aldrich®
Ethidium bromide solution 0,025 % in dropper bottle	aqueous solution (250 µg/ml)
Ethylene glycol	for analysis EMSURE® Reag. Ph Eur, Reag. USP.
Ferric ammonium citrate	ca. 18 % Fe, DAC; Microbiology
Ficoll 400	BioXtra, for molecular biology, lyophilized powder
Formaldehyde solution	Formaldehydlösung 37 % ≥37 %, Ph.Eur., BP, USP
Formic acid 98%	ROTIPURAN® ≥98 %, p.a., ACS
Gallium	99.9995% Metals basis
Gentamycin sulphate	≥590 I.U./mg, for biochemistry
Glutaraldehyde solution	grade II, 25% in H ₂ O
Glutaraldehyde solution, 8%	Grade I, 8% in H ₂ O, specially purified for use as an electron microscopy fixative
Glycerol / Glycerin	ROTIPURAN® ≥99,5 %, p.a., anhydrous
Glycerol / Glycerin	ROTIPURAN
Glycid ether 100	for electron microscopy
Glycine	PUFFERAN® ≥99 %, p.a.
Guanidine hydrochloride	≥99,5 %, for biochemistry
Guanosine 5'-triphosphate sodium salt hydrate	≥95% (HPLC), powder
Hellmanex III	
HEPES	PUFFERAN® ≥99,5 %, p.a. For biochemistry and molecular biology.
HEPES sodium salt	
Heptane	reagent grade, 99%
Hexane	puriss. p.a., ACS reagent, reagent Ph. Eur., ≥99% (GC)
Hydrochloric acid	37% EMSURE® ACS, ISO, Reagent Ph. Eur. analytical reagent
Hydrochloric acid	37% in water analytical reagent
Hydrochloric Acid 32%	32 %, extra pure
Hydrogen peroxide 30%	Ph.Eur., stabilized
Hydrogenperoxid	33%
Hypophosphorous acid solution 50 wt. % in H ₂ O	
Imidazole	PUFFERAN® ≥99 %, p.a., ultra quality
Imidazole	PUFFERAN® ≥99 %, p.a., ultra quality
Immersion oil	for microscopy
Insulin from bovine pancreas	powder, BioReagent, suitable for cell culture
Iotect - iod indicator	Starch, soluble
IPTG	Isopropyl-β-D-thiogalactopyranosid
IPTG	≥99 %, BioScience-Grade, dioxane free, animal free
IPTG	≥99 %, BioScience-Grade, dioxane free, animal free
Iron (II) sulphate heptahydrate	nalaR NORMAPUR® Reagent Ph. Eur. analytical reagent
Iron(II) sulfate heptahydrate	for analysis EMSURE® ACS, ISO, Reagent Ph Eur.
Iron(III) chloride hexahydrate	reagent grade, ≥98%
Isoamyl alcohol	EMSURE® ACS, Reagent Ph. Eur. zur Analyse

Isopropanol	hypergrade for LC-MS LiChrosolv, 99%
Isopropanol	LC-MS CHROMASOLV, 99%
Isopropanol	HiPerSolv CHROMANORM, 99%
Kanamycin sulphate	≥750 I.U./mg, for biochemistry
L(-)-Malic acid	≥99 %, for biochemistry
L-(+)-Arabinose	≥99%
Lactic acid	≥85%
L-Alanine	BioUltra, ≥99.5% (NT)
L-Arginine	≥98,5 %, USP, for biochemistry
L-Asparagine	≥98% (HPLC)
L-Aspartic acid sodium salt monohydrate	≥98% (TLC)
LB Broth (Lennox)	for molecular biology
L-Cysteine	BioUltra, ≥98.5% (RT)
L-Glutamic acid	from non-animal source, meets EP testing specifications, suitable for cell culture, 98.5-100.5%
L-glutamic acid monopotassium salt monohydrate	
L-Glutamic acid monosodium salt; Natrium-L(+)-glutamat Monohydrat	
L-Glutamine	<i>ReagentPlus</i> [®] , ≥99% (HPLC)
L-glutathione reduced	>98 %, for biochemistry γ-L-Glutamyl-L-cysteinyl-glycine, GSH
L-Histidine	cell culture tested, meets EP, USP testing specifications, from non-animal source
Lincomycin	Lincomycin hydrochloride
Lipase	Lipase, <i>Chromobacterium viscosum</i>
Lipase, <i>Chromobacterium viscosum</i>	
Liquified phenol	meets USP testing specifications, ≥89.0%
L-Isoleucine	CELLPURE [®] ≥99 %
Lithium acetate dihydrate	BioXtra
Lithium chloride	≥99,5 %, p.a., ultra quality
Lithium nitrate	≥99,5 %, anhydrous
Lithium sulphate	≥99 %, anhydrous
L-Leucine	from non-animal source, meets EP, JP, USP testing specifications, suitable for cell culture, 98.5-101.0%
L-Lysine hydrochloride	CELLPURE [®] ≥99 %
L-Lysine monohydrochloride	from non-animal source, meets EP, JP, USP testing specifications, suitable for cell culture, 98.5-101.0%
L-Methionine nonanimal source	from non-animal source, meets EP, JP, USP testing specifications, suitable for cell culture, 99.0-101.0%
L-Phenylalanine	reagent grade, ≥98%
L-Proline	<i>ReagentPlus</i> [®] , ≥99% (HPLC)
L-Seleno-Methionine	
L-Serine	from non-animal source, meets EP, USP testing specifications, suitable for cell culture, 98.5-101.0%
L-Threonine	reagent grade, ≥98% (HPLC)
L-Tryptophane	BioUltra, ≥99.5% (NT)
L-Tyrosine	reagent grade, ≥98% (HPLC)

L-Valine	from non-animal source, meets EP, JP, USP testing specifications, suitable for cell culture, 98.5-101.0%
Lysozyme	lyophilized powder, protein ≥90 %, ≥40,000 units/mg protein
Lysozyme	≥35000 FIP U/mg, BioScience-Grade, lyophilised
Lysozyme	≥35000 FIP U/mg, BioScience-Grade, lyophilised
Magnesium acetate tetrahydrate	BioXtra, ≥99%
Magnesium chloride	CELLPURE® ≥99 % For cell culture and biochemistry.
Magnesium chloride hexahydrate	≥99 %, p.a., ACS
Magnesium chloride hexahydrate	
Magnesium formate dihydrate	BioXtra, ≥98.0% (RT)
Magnesium nitrate hexahydrate	≥98 %, p.a., ACS
Magnesium sulfate heptahydrate	≥99 %, Ph.Eur., USP, BP
Magnesium sulphate hydrate	≥99 %, dried, extra pure, DAC
Maleic acid	<i>ReagentPlus</i> ®, ≥99.0% (HPLC)
Malonic acid	for synthesis
Mangan(II)sulfat	Monohydrat, >98%
Manganese(II)chloride tetrahydrate	≥99 %, p.a.
MES	PUFFERAN® ≥99 %
Methanol	≥99,9 %
Methanol	ROTIPURAN® ≥99,9 %, p.a., ACS, ISO
Methanol	>99,9%, spectrophotometric grade
Methanol	>99,9%, extra dry
Methanol	>99,9%, extra dry
Methylnadic anhydride	pract.
Milk powder	blotting grade, powdered, low fat
MOPS	PUFFERAN® ≥99,5 %
N,N-Dimethylformamid	
N,N-Dimethylformamide	for molecular biology, ≥99%
n-Decane	≥95%
n-Dodecyl β-maltoside	
n-Heptyl β-D-thioglucopyranoside	≥99%
n-Hexane	Emplura
Nickel (II) Sulphate hexahydrate	≥99 %, p.a., ACS
Nickel(II)chloride hexahydrate	BioReagent, suitable for cell culture
Nitric acid	65%
Oil for heating bath	Rotitherm K+H
ortho-Phosphoric acid 85 %	ROTIPURAN® ≥85 %, p.a., ISO
Paraformaldehyde	prilled, 95%
Paraformaldehyde, in vials	16% formaldehyde, in vials, 10 x 10 mL
PEG ME 5000	Poly(ethylene glycol) methyl ether average Mn 5,000
PENICILLIN -STREPTOMYCIN	
Penicillin-Streptomycin	
Pepsin from porcine gastric mucosa	lyophilized powder, 3,200-4,500 units/mg protein
Peptone ex casein	tryptic digest, for microbiology
Perfluorodecalin	95%

Petroleum ether, low boiling point	30 - 40 °C 99.5-101.0%, puriss. zur Analyse, Riedel-de Haën™
pH buffer for pH meter calibration	pH 4.01
pH buffer for pH meter calibration	pH 7.01
pH buffer for pH meter calibration	pH 10.1
Phenol/Chloroform/Isoamylalcohol	Roti®-Phenol/Chloroform/Isoamylalcohol
Phosphoric acid	85%
PIPES	PUFFERAN® ≥99 %
Poly(ethylene glycol) 200, PEG 200	BioUltra, 200
Poly(ethylene glycol) 400, PEG 400	BioUltra, 400
Polyethylene glycol (PEG) 1000	BioUltra, 1,000
Polyethylene glycol (PEG) 10000	BioUltra, 10,000
Polyethylene glycol (PEG) 12000	12,000
Polyethylene glycol (PEG) 1500	BioUltra, 1,500
Polyethylene glycol (PEG) 2000	ROTIPURAN® Ph.Eur.
Polyethylene glycol (PEG) 3000	BioUltra, 3,000
Polyethylene glycol (PEG) 3350	BioUltra, 3,350
Polyethylene glycol (PEG) 4000	
Polyethylene glycol (PEG) 6000	BioUltra, for molecular biology, 6,000
Polyethylene glycol (PEG) 8000	BioUltra, for molecular biology, 8,000
Ponceau S solution	BioReagent, suitable for electrophoresis, 0.1 % (w/v) in 5% acetic acid
Potassium acetate	≥99 %, Ph.Eur., BP
Potassium bromate	≥99.8%, EMSURE® ACS, ISO, Reag. Ph. Eur. zur Analyse
Potassium bromide	Uvasol® for IR spectrophotometry, Supelco®
Potassium chloride	
Potassium chloride	≥99,5 %, p.a., ACS, ISO
Potassium dihydrogen phosphate	>99%, p.a., ACS
Potassium dihydrogen phosphate	>99%, p.a., ACS
Potassium dihydrogen phosphate	>99%, p.a., ACS
Potassium dihydrogen phosphate	>99%, p.a., ACS
Potassium dihydrogen phosphate	>99%, p.a., ACS
Potassium ferricyanide; Potassium hexacyanoferrate(III)	ACS reagent, ≥99.0%
Potassium formate	ReagentPlus®, 99%
Potassium hydrogen sulphate	≥99 %, extra pure
Potassium Hydroxide Pellets	≥85 %, Ph.Eur., pellets
Potassium hydroxide solution 50%	Kalilauge 50 % reinst - 1 l
Potassium iodate	99,7-100,4%
Potassium iodide	Emsure
Potassium nitrate	≥99 %, Ph.Eur., BP
Potassium phosphate monobasic	ReagentPlus®
Potassium sulphate	≥99 %, p.a., ACS, ISO
Propylene glycol monomethyl ether acetate (PGMEA)	≥99.5%
PROTAMINE SULFATE	
Proteinase K	lyophil., ≥30 mAnson U/mg
Pyridine	anhydrous (max. 0.003% H ₂ O) ≥99.8%
SDS Pellets	≥99 %, for biochemistry
Silica Gel with moisture indicator (brwon gel)	desiccant ~ 1 - 4 mm

Silicone Oil	DC200 Silicone Oil
Silicone solution SERVA	in isopropanol
SOB Broth	for molecular biology
Sodium acetate	≥99 %, p.a., ACS, anhydrous Dehydrating agent, for molecular biology
Sodium acetate	anhydrous, BioUltra, for luminescence, for molecular biology, ≥99.0% (NT)
Sodium acetate trihydrate	≥99,5 %, p.a., ACS, ISO
Sodium bicarbonate	puriss. p.a., ACS reagent, reag. Ph. Eur., ≥99.7%, powder
Sodium bromide	≥99 %, p.a., ACS
Sodium chloride	≥99,5 %, p.a., ACS, ISO
Sodium chloride	≥99,5 %, p.a., ACS, ISO
Sodium chloride	≥99,5 %, p.a., ACS, ISO For freezing mixtures; salting-out soap. For molecular biology.
Sodium citrate tribasic dihydrate	for molecular biology, ≥99%
Sodium citrate tribasic dihydrate	ACS reagent, ≥99.0%
Sodium dihydrogen phosphate (*2H2O)	Dihydrate for analysis EMSURE® Reag. Ph Eur 1
Sodium dihydrogen phosphate dihydrate	≥99 %, p.a. For molecular biology.
Sodium dihydrogen phosphate monohydrate	≥98 %, p.a., ACS. Molecular biology.
Sodium disulfite	Sodium disulfite (sodium metabisulfite) for analysis EMSURE® ACS,Reag. Ph Eur.
Sodium dithionite	
Sodium fluoride	≥99 %, p.a., ACS, ISO
Sodium formate	≥99 %, p.a., ACS
Sodium hydroxide	
Sodium hydroxide	10% solution
Sodium hydroxide solution 20%	Natronlauge 20 % reinst - 2,5 l
Sodium hydroxide, pellets	≥99 %, p.a., ISO, in pellets
Sodium hypochlorite solution	12 % Cl, techn.
Sodium iodide	≥99 %, Ph.Eur.
Sodium malonate	Malonic acid disodium monohydrat
Sodium nitrate	BioXtra, ≥99.0%
Sodium nitrite	≥99.0% AnaLaR NORMAPUR® for Analyse
Sodium sulphate	≥99 %, p.a., ACS, ISO, powdered, anhydrous
Sodium thiocyanate	≥98 %, p.a., ACS
Sodium thiosulphate	EMPLURA
Spectinomycin	Spectinomycin sulfate, 93%
Spectinomycin dihydrochloride pentahydrate	potency: ≥603 µg per mg
Streptomycin sulphate	>720 I.U./mg, for biochemistry
Succinic acid	≥99 %, p.a., ACS
Sulphuric acid	ROTIPURAN® 98%
Sulphuric acid 25 %	25 %, extra pure
TBE Buffer 10x	Rotiphorese® 10x TBE Buffer
TechniEtch Cr01	N/A
TEMED	≥99 %, p.a., for electrophoresis

Tetracycline Hydrochloride	CELLPURE® ≥900 µg/mg
Tetrahydrofurane	ACROS, Extra dry
Tetrahydrofurane	ACROS, Extra dry
Tetrahydrofurane	Tetrahydrofuran ROTISOLV® HPL-1 liter
TETRAMETHYLAMMONIUMBROMID	
TGX Stain-Free FastCast Acrylamide, 12%, 50-Gel Kit	
Thaumatococcus daniellii	
Thiamine hydrochloride	
Toluol	ROTIPURAN® ≥99,5 %, p.a., ACS, ISO
Toluol	<i>anhydrous (max. 0.005% H₂O), SeccoSolv®, Sigma-Aldrich®</i>
Toluol	<i>anhydrous (max. 0.005% H₂O) ≥99.8%, AnalaR NORMAPUR® analytical reagen</i>
TRICINE	PUFFERAN® ≥99 %
TRIS	PUFFERAN® ≥99,9 %, ultra quality
TRIS	PUFFERAN® ≥99,3 %, Buffer Grade
TRIS hydrochloride	PUFFERAN® ≥99 %, p.a.
Tris-(2-carboxyethyl)-phosphine hydrochloride	≥98 %, for biochemistry
tri-Sodium phosphate dodecahydrate	≥98 %, p.a., ACS
Triton® X-100	laboratory grade
Trypsin from bovine pancreas	Type I, ~10,000 BAEE units/mg protein
Trypsin-Inhibitor	≥10000 BAEE U/mg
Tryptone/Peptone ex casein	pancreatic digest, for microbiology
Tryptone/Peptone ex casein	pancreatic digest, for microbiology
TWEEN 80	Ph.Eur. Non-ionic detergent.
Tween® 20	Ph.Eur. Non-ionic detergent.
Urea	≥99,5 %, p.a. For microbiology and molecular biology.
Urea	Harnstoff min. 99,5 %, p.a.
Water	BioScience Grade, nuclease-free, steam sterilised, DEPC treated water
X-beta-Gal	
Xylene cyanol (C.I. 42135)	for histology and electrophoresis
Xylose	D-Xylose
Xylose	D-Xylose
Yeast Extract	powdered, for bacteriology
Zinc acetate dihydrate	≥99,5 %, p.a., ACS
Zinc chloride	≥97 %, p.a.
Zinc sulfate heptahydrate	puriss. p.a., ACS reagent, reag. ISO, reag. Ph. Eur., ≥99.5%